

Type specimens of the Apocynaceae, Asclepiadaceae, Gentianaceae, Loganiaceae and Menyanthaceae in the Herbarium of the Martin-Luther-University Halle Wittenberg (HAL)

STEFAN KESTING & MATTHIAS H. HOFFMANN

Abstract: KESTING, S. & HOFFMANN, M. H. 2000: Type specimens of the Apocynaceae, Asclepiadaceae, Gentianaceae, Loganiaceae and Menyanthaceae in the Herbarium of the Martin-Luther-University Halle Wittenberg. *Schlechtendalia* 5: 11-17.

Asclepiadaceae

Astephanus mucronatus Schldl., *Linnaea* 8: 518, 1833.

"N. 10 ... Hac. de la Laguna Jul. 29. ... [written by C.J.W. Schiede]" (holotype: HAL 82965, 2 sheets).

Sarcostemma brevistigma Wight & Arn., in Wight, *Contrib.* 59, 1834.

"Herb. Wight. propr. 1558. *Sarcostemma brevistigma* W. et Arn. Peninsula Ind. orientalis." (type: HAL 83639).

Tacazzea venosa Decne., DC. *Prod.* 8: 493, 1844.

"Schimperiter Abyssinicum ... 636. *Periploca venosa* Hochst. Ad ripas fluvii Tacaze infra Dscheladscheranne d. 8. Nov. 1839." (type: HAL 83637).

Toxocarpus kleinii Wight & Arn., *Contrib.* 61, 1834.

"Herb. Wight. propr. 1560. *Toxocarpus Kleinii* W. et Arn. Peninsula Ind. orientalis." (type: HAL 83638).

Tylophora koi Merrill, *Sunyatsenia* 2: 17, 1934.

"*Tylophora koi* Merrill, sp. nov., S.P. Ko 51732, W.Y. Chun (otherwise written in Chinese)" (isotype: HAL 85404).

Original material of the Asclepiadaceae

Periploca linearis Hochst., *Flora* 24, 1. *Intell.*: 25, 1841, nom. nud.

"Schimperiter Abyssinicum ... 365. *Periploca linearis* Hochst. ... In regione media septentrionali montis Scholoda frutices et arbores caule volubili scandens d. 28. Oct. 1837." (HAL 83636).